Ernemann, U.
—, Schirmer, D., Wöber, A., Langmann, P., Warzecha, K., Joeres, R., Klinker, H. and Richter, E.
Gas chromatographic determination of trimethoprim in plasma during high-dose trimethoprim–sulfamethoxazole treatment for *Pneuymocystis carinii* pneumonia 243
Ersser, R.S., see Davey, J.F. 9
Espinosa, J., see Ladrón De Guevara, O. 35
Eto, K., see Gomita, Y. 509
Evans, P.C., see Young, D. 385
Ferry, S., see Boulieu, R. 542
Fleming, R.A.
— and Stewart, C.F.
Stability-indicating high-performance chromatographic method for the simultaneous determination of cisplatin and 5-fluorouracil in 0.9% sodium chloride for injection 517
Flores-Murietta, F.J.
—, Castañeda-Hernández, G. and Hong, E.
Determination of pelanserin in human plasma by high-performance liquid chromatography with amperometric detection 501
Fuhrer, J.P.
Rapid analysis of membrane glycopeptides by gel permeation high-performance liquid chromatography 65
Fujimori, H.
—, Sasaki, T., Hibi, K., Senda, M. and Yoshioka, M.
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305
Fujita, K., see Giegerich, G. 79
Fukuda, T., see Gomita, Y. 509
Furuno, K., see Gomita, Y. 509
Gärtner, W., see Dudot, C. 435
Gaitonde, C.D.
— and Pathak, P.V.
Estimation of the analgesic muscle relaxants chlorzoxazone and diazepam in human plasma by reversed-phase liquid chromatography 407
García, L., see Ladrón De Guevara, O. 35
Giegerich, G.
—, Pette, M., Fujita, K., Wekerle, H., Epplen, J.T. and Hinkkanen, A.
Rapid method based on reversed-phase high-performance liquid chromatography for purification of human myelin basic protein and its thrombic and endoproteinase Lys-C-peptides 79
Girardot, R., see Baldacini, O. 357
Goldberg, V.D., see Elyas, A.A. 473
Gomita, Y.
—, Furuno, K., Eto, K., Fukuda, T., Araki, Y., Yamori, M. and Oishi, R.
Rapid and simple determination of nicorandil in rat plasma using a solid-phase extraction column 509
Gordon, E.R., see Goresky, C.A. 123
Gorecki, D.K.J.
—, Hindmarsh, K.W., Hall, C.A., Mayers, D.J. and Sankaran, K.
Determination of chloral hydrate metabolism in adult and neonate biological fluids after single-dose administration 333
Goresky, C.A
— and Gordon, E.R.
High-performance liquid chromatographic separation of bilirubin conjugates: the effects of change in molarity and pH 123
Green, G.A., see Baldacini, O. 357
Greig, N.H., see Deutsch, J. 325
Guelen, P.J.M., see Lansdorp, D. 487
Hall, C.A., see Gorecki, D.K.J. 333
Hall, S.D., see Rudy, A.C. 395
Hayashi, T.
—, Todoriki, H. and Iida, Y.
Highly sensitive method for the determination of 1-methyl-1,2,3,4-tetrahydro-$\beta$-carboline using combined capillary gas chromatography and negative-ion chemical ionization mass spectrometry 1
Hayball, P.J.
—, Cosh, D.G. and Wrobel, J.
Rapid, sensitive determination of human serum midazolam by high-performance liquid chromatography 526
Hellgren, U.
—, Villén, T. and Ericsson, Ö.
High-performance liquid chromatographic determination of quinine in plasma, whole blood and samples dried on filter paper 221
Hibi, K., see Fujimori, H. 305
Higashiyama, K., see Kataoka, H. 172

Higuchi, S., see Teramura, T. 191
Hindmarsh, K.W., see Gorecki, D.K.J. 333
Hinkkanen, A., see Giegerich, G. 79
Hiro, T.
—, Kadofuku, T., Makino, Y. and Sato, T.
Detection of the exercise-associated changes in serum proteins by two-dimensional electrophoresis under non-denaturing conditions 315
Hirose, S., see Kito, M. 91
Holden, C.L., see Laizure, S.C. 235
Hong, E., see Flores-Murietta, F.J. 501
Iida, Y., see Hayashi, T. 1
Jaenisch, R., see Ruprecht, R.M. 371
Jakubowski, E.M.
—, Woodard, C.L., Mershon, M.M. and Dolzine, T.W.
Quantification of thiodiglycol in urine by electron ionization gas chromatography-mass spectrometry 184
Janssen, T.J., see Lansdorp, D. 487
Jarvi, E.J., see Patrick, K.S. 214
Joeres, R., see Ernemann, U. 243
Johnston, A., see Saleh, S. 531
Jurado, M.C., see Martínez, D. 459
Kadofuku, T., see Hiro, T. 315
Kahn, R.S., see Suckow, R.F. 228
Kanazawa, T., see Yamano, Y. 199
Kandiko, C.T.
—, Browning, S., Cooper, T. and Cox, W.A.
Detection of low nanogram quantities of phencyclidine extracted from human urine. Preparation of an acetylated column packing material for use in gas chromatography with nitrogen–phosphorus detection 208
Kataoka, H.
—, Higashiyama, K., Okazaki, T. and Makita, M.
Determination of isethionic acid by gas chromatography with flame photometric detection 172
Kazemifard, A.G.
— and Morgan, D.J.
Determination of busulfan in plasma by high-performance liquid chromatography 274
Keizer, H., see Roemen, T.H.M. 447
Kelly, M.T.
—, Smyth, M.R. and Ruskin, H.
Evaluation of drug-free plasma profiles by high-performance liquid chromatography following on-line solid-phase extraction 277
Kito, M.
—, Tawa, R., Takeshima, S. and Hirose, S.
Determination of purine nucleosides and their bases by high-performance liquid chromatography using co-immobilized enzyme reactors 91
Klein, C., see Dudot, C. 435
Klinker, H., see Ernemann, U. 243
Kobayashi, S., see Teramura, T. 191
Krämer, B.K., see Kühlkamp, V. 267
Kühlkamp, V.
—, Schmid, F., Ress, K.M., Krämer, B.K., Mayer, F., Liebich, H.M., Risler, T. and Seipel, L.
Quantification of cibenzoline and its imidazole metabolite by high-performance liquid chromatography in human serum 267
Kwon, K.I., see Lee, H.S. 425
Lachatre, G., see Nicot, G. 258
Ladrón De Guevara, O.
—, Cortinas De Nava, O.C., Padilla, P., Espinosa, J., Cebrian, M. and García, L.
Rapid determination of β-aminoisobutyric acid by reversed-phase high-performance liquid chromatography 35
Laizure, S.C.
—, Holden, C.L. and Stevens, R.C.
Ion-paired high-performance liquid chromatographic separation of trimethoprim, sulfamethoxazole and $N^4$-acetylsulfamethoxazole with solid-phase extraction 235
Lajdová, I.
—, Sputová, V., Mikula, J., Černay, P. and Dzúrik, R.
Isolation of an additional inhibitor of glucose utilization in renal insufficiency: pseudouridine 178
Langguth, P.
—, Spahn, H. and Merkle, H.-P.
Fluorescence assay for small peptides and amino acids: high-performance liquid chromatographic determination of selected substrates using activated *S*-flunoxaprofen as a chiral derivatizing agent 55
Langmann, P., see Ernemann, U. 243

Lansdorp, D.
—, Janssen, T.J., Guelen, P.J.M. and Vree, T.B.
High-performance liquid chromatographic method for the determination of diclofenac and its hydroxy metabolites in human plasma and urine 487
Lee, H.S.
—, Zee, O.P. and Kwon, K.I.
Simultaneous determination of cefamandole and cefamandole nafate in human plasma and urine by high-performance liquid chromatography with column switching 425
Lee Chin, J.R.
Determination of six indolic compounds, including melatonin, in rat pineal using high-performance liquid chromatography with serial fluorimetric–electrochemical detection 111
Leenheer, A.P. De, see Van Liederkerke, B.M. 155
Liebich, H.M., see Kühlkamp, V. 267
Liederkerke, B.M. Van, see Van Liederkerke, B.M. 155
Makino, Y., see Hiro, T. 315
Makita, M., see Kataoka, H. 172
Marquet, P., see Nicot, G. 258
Martínez, D.
—, Jurado, M.C. and Repetto, M.
Analysis of buprenorphine in plasma and urine by gas chromatography 459
Matheson, L.M., see Cummings, J. 43
Mayer, F., see Kühlkamp, V. 267
Mayers, D.J., see Gorecki, D.K.J. 333
Mazrimas, J.A., see Stacks, P.C. 25
Mazzi, G.
— and Schinella, M.
Simple and practical high-performance liquid chromatographic assay of propofol in human blood by phenyl column chromatography with electrochemical detection 537
Merkle, H.-P., see Langguth, P. 55
Mershon, M.M., see Jakubowski, E.M. 184
Meulemans, A., see Mohler, J. 415
Mikula, J., see Lajdová, I. 178
Mohler, J.
—, Meulemans, A. and Vulpillat, M.
High-performance liquid chromatographic determination of the binding of ceftriaxone to human serum albumin solution and albumin from diluted human serum 415
Monteil, H., see Baldacini, O. 357
Moreau, J., see Dudot, C. 435
Morgan, D.J., see Kazemifard, A.G. 274
Morishita, N., see Yamano, Y. 199
Nakai, H., see Yamano, Y. 199
Nicot, G.
—, Lachatre, G., Marquet, P., Bonnaud, F., Vallette, J.P. and Rocca, J.-L.
High-performance liquid chromatographic determination of navelbine in human plasma and urine 258
Nill, K., see Nusser, E. 163
Nusser, E.
—, Nill, K. and Breyer-Pfaff, U.
Enantiomer analysis of *E*- and *Z*-10-hydroxyamitriptyline in human urine 163
Ogawa, T., see Yamano, Y. 199
Oishi, R., see Gomita, Y. 509
Okazaki, T., see Kataoka, H. 172
Ostrowski, J., see Dudot, C. 435
Padilla, P., see Ladrón De Guevara, O. 35
Pathak, P.V., see Gaitonde, C.D. 407
Patrick, K.S.
— and Jarvi, E.J.
Capillary gas chromatographic–mass spectrometric analysis of plasma methylphenidate 214
Patsalos, P.N., see Elyas, A.A. 473
Paxton, J.W., see Young, D. 385
Pays, M., See Dudot, C. 435
Pette, M., see Giegerich, G. 79
Rapoport, S.I., see Deutsch, J. 325
Repetto, M., see Martínez, D. 459
Ress, K.M., see Kühlkamp, V. 267
Richards, D.A., see Sherwood, R.A. 293
Richter, E., see Ernemann, U. 243
Rihn, B., see Baldacini, O. 357
Risler, T., see Kühlkamp, V. 267
Rocca, J.-L., see Nicot, G. 258
Rodriguez, M., see Alfonso, D. 101
Roemen, T.H.M.
—, Keizer, H. and Van der Vusse, G.J.
Quantitative assessment of γ-linolenic acid in human blood and plasma with capillary gas chromatography 447
Roosemalen, M.C.M., see Soons, P.A. 343
Rudy, A.C.
—, Anliker, K.S. and Hall, S.D.

High-performance liquid chromatographic determination of the stereoisomeric metabolites of ibuprofen 395
Ruprecht, R.M.
—, Sharpe, A.H., Jaenisch, R. and Trites, D.
Analysis of 3′-azido-3′-deoxythymidine levels in tissues and milk by isocratic high-performance liquid chromatography 371
Ruskin, H., see Kelly, M.T. 277
Rustum, A.M.
Measurement of verapamil in human plasma by reversed-phase high-performance liquid chromatography using a short octyl column 480
Saleh, S.
—, Johnston, A. and Turner, P.
Measurement of medifoxamine metabolites in urine by high-performance liquid chromatography 531
Samuelsen, O.B.
High-performance liquid chromatographic determination of furazolidone in Atlantic salmon (*Salmo salar*) tissue 495
Sankaran, K., see Gorecki, D.K.J., 333
Sasaki, T., see Fujimori, H. 305
Sato, T., see Hiro, T. 315
Schinella, M., see Mazzi, G. 537
Schirmer, D., see Ernemann, U. 243
Schmid, F., see Kühlkamp, V. 267
Seipel, L., see Kühlkamp, V. 267
Senda, M., see Fujimori, H. 305
Sharpe, A.H., see Ruprecht, R.M. 371
Shayman, J.A.
— and Barcelon, F.S.
Ion-pair chromatography of inositol polyphosphates with N-methylimipramine 143
Sherwood, R.A.
—, Titheradge, A.C. and Richards, D.A.
Measurement of plasma and urine amino acids by high-performance liquid chromatography with electrochemical detection using phenylisothiocyanate derivatization 293
Smyth, J.F., see Cummings, J. 43
Smyth, M.R., see Kelly, M.T. 277
Soncrant, T.T., see Deutsch, J. 325
Soons, P.A.
—, Roosemalen, M.C.M. and Breimer, D.D.
Enantioselective determination of felodipine and other chiral dihydropyridine calcium entry blockers in human plasma 343
Spahn, H., see Langguth, P. 55
Spreux-Varoquaux, O., see Dudot, C. 435
Spustová, V., see Lajdová, I. 178
Stacks, P.C.
—, Mazrimas, J.A., Corzett, M. and Balhorn, R.
Reactivity and adduct formation of a polyaromatic hydrocarbon, 7-bromomethylbenz[*a*]anthracene, with chromatin histone proteins 25
Stevens, R.C., see Laizure, S.C. 235
Stewart, C.F., see Fleming, R.A. 517
Suckow. R.F.
—, Cooper, T.B. and Kahn, R.S.
High-performance liquid chromatographic method for the analysis of plasma *m*-chlorophenylpiperazine 228
Sudo, A.
Analysis of corticosterone in rat urine by high-performance liquid chromatography and fluorimetry using post-column reaction with sulphuric acid 453
Swart, P.J.
—, Drenth, B.F.H. and De Zeeuw, R.A.
Reversed-phase liquid chromatographic method with amperometric detection for the determination of the dopamine agonist 2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin (N-0437) in human plasma and urine 464
Takeshima, S., see Kito, M. 91
Tawa, R., see Kito, M. 91
Teramura, T.
—, Kobayashi, S. and Higuchi, S.
Determination of mepirodipine in plasma by capillary column gas chromatography-negative-ion chemical ionization mass spectrometry 191
Titheradge, A.C., see Sherwood, R.A. 293
Todoriki, H., see Hayashi, T. 1
Trites, D., see Ruprecht, R.M. 371
Turner, P., see Saleh, S. 531
Vallette, J.P., see Nicot, G. 258
Van der Vusse, G.J., see Roemen, T.H.M. 447
Van Liederkerke, B.M.
— and De Leenheer, A.P.
Analysis of xanthene dyes by reversed-

phase high-performance liquid chromatography on a polymeric column followed by characterization with a diode array detection 155
Villén, T., see Hellgren, U. 221
Vree, T.B., see Lansdorp, D. 487
Vulpillat, M., see Mohler, J. 415
Vusse, G.J. van der, see Roemen, T.H.M. 447
Warzecha, K., see Ernemann, U. 243
Wekerle, H., see Giegerich, G. 79
Wöber, A., see Ernemann, U. 243
Woodard, C.L., see Jakubowski, E.M. 184
Wrobel, J., see Hayball, P.J. 526
Yamada, K., see Yamano, Y. 199
Yamaghishi, Y., see Yamano, Y. 199
Yamano, Y.
—, Nakai, H., Ogawa, T., Kanazawa, T., Morishita, N., Yamada, K. and Yamagishi, Y. Determination of 4-cyano-5,5-bis(4-methoxyphenyl)-4-pentenoic acid in human plasma and platelets by gas chromatography–mass spectrometry 199
Yamori, M., see Gomita, Y. 509
Yoshioka, M., see Fujimori, H. 305
Young, D.
—, Evans, P.C. and Paxton, J.W. Quantitation of the antitumour agent N-[2-(dimethylamino)ethyl]acridine-4-carboxamide in plasma by high-performance liquid chromatography 385
Zee, O.P., see Lee, H.S. 425
Zeeuw, R.A. de, see Swart, P.J. 464

# Subject Index


Acetaminophen
Estimation of the analgesic muscle relaxants chlorzoxazone and diazepam in human plasma by reversed-phase LC 407

N-Acetylserotonin
Determination of six indolic compounds, including melatonin, in rat pineal using HPLC with serial fluorimetric–electrochemical detection 111

$N^4$-Acetylsulfamethoxazole
Ion-paired HPLC separation of trimethoprim, sulfamethoxazole and $N^4$-acetylsulfamethoxazole with solid-phase extraction 235

Adenine nucleotides
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305

Adenosine
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305

ADP
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305

Albumin
HPLC determination of the binding of ceftriaxone to human serum albumin solution and albumin from diluted human serum 415

Amino acids
Fluorescence assay for small peptides and amino acids: HPLC determination of selected substrates using activated *S*-flunoxaprofen as a chiral derivatizing agent 55

Amino acids
Measurement of plasma and urine amino acids by HPLC with electrochemical detection using phenylisothiocyanate derivatization 293

Amino acids, PTC derivatives
Amino acid analysis of physiological fluids by HPLC with phenylisothiocyanate derivatization and comparison with ion-exchange chromatography 9

$\beta$-Aminoisobutyric acid
Rapid determination of $\beta$-aminoisobutyric acid by reversed-phased HPLC 35

AMP
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305

ATP
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305

3′-Azido-3′-deoxythymidine
Analysis of 3′-azido-3′-deoxythymidine levels in tissues and milk by isocratic HPLC 371

Bilirubin conjugates
HPLC separation of bilirubin conjugates: the effects of change in molarity and pH 123

7-Bromomethylbenz[*a*]anthracene
Reactivity and adduct formation of a polyaromatic hydrocarbon, 7-bromomethylbenz[*a*]anthracene, with chromatin histone proteins 25

Buprenorphine
Analysis of buprenorphine in plasma and urine by GC 459

Busulfan
Determination of busulfan in plasma by HPLC 274

cAMP
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305

Cefamandole
Simultaneous determination of cefamandole and cefamandole nafate in human plasma and urine by HPLC with column switching 425

Cefamandole nafate
Simultaneous determination of cefamandole and cefamandole nafate in human plasma and urine by HPLC with column switching 425

Ceftriaxone
HPLC determination of the binding of ceftriaxone to human serum albumin solution and albumin from diluted human serum 415

Chloral hydrate
Determination of chloral hydrate metabolism in adult and neonate biological fluids after single-dose administration 333
*m*-Chlorophenylpiperazine
HPLC method for the analysis of plasma *m*-chlorophenylpiperazine 228
Chlorzoxazone
Estimation of the analgesic muscle relaxants chlorzoxazone and diazepam in human plasma by reversed-phase LC 407
Cibenzoline
Quantification of cibenzoline and its imidazole metabolite by HPLC in human serum 267
Cisplatin
Stability-indicating HPLC method for the simultaneous determination of cisplatin and 5-fluorouracil in 0.9% sodium chloride for injection 516
*Clostridium sordellii* cytotoxin
Fast purification of *Clostridium sordellii* cytotoxin 357
Corticosterone
Analysis of corticosterone in rat urine by HPLC and fluorimetry using post-column reaction with sulphuric acid 453
4-Cyano-5,5-bis(4-methoxyphenyl)-4-pentenoic acid
Determination of 4-cyano-5,5-bis(4-methoxyphenyl)-4-pentenoic acid in human plasma and platelets by GC–MS 199
Cytotoxins
Fast purification of *Clostridium sordellii* cytotoxin 357
N-Demethyldeacetyldiltiazem
Solid-phase extraction of diltiazem and its metabolites from plasma prior to HPLC 542
N-Demethyldiltiazem
Solid-phase extraction of diltiazem and its metabolites from plasma prior to HPLC 542
Desacetyldiltiazem
Solid-phase extraction of diltiazem and its metabolites from plasma prior to HPLC 542
N-Desmethylmedifoxamine
Measurement of medifoxamine metabolites in urine by HPLC 531
Diazepam
Estimation of the analgesic muscle relaxants chlorozoxazone and diazepam in human plasma by reversed-phase LC 407
Diclofenac
HPLC method for the determination of diclofenac and its hydroxy metabolites in human plasma and urine 487
4,5-Dihydro-2-(2,2-diphenylcyclopropyl)-1*H*-imidazole
Quantification of cibenzoline and its imidazole metabolite by HPLC in human serum 267
Dihydropyridine calcium entry blockers
Enantioselective determination of felodipine and other chiral dihydropyridine calcium entry blockers in human plasma 343
3,4-Dihydroxyphenylacetic acid
Method for the determination of γ-L-glutamyl-L-dihydroxyphenylalanine and its major metabolites L-dihydroxyphenylalanine, dopamine and 3,4-dihydroxyphenylacetic acid by HPLC with electrochemical detection 43
L-Dihydroxyphenylalanine
Method for the determination of γ-L-glutamyl-L-dihydroxyphenylalanine and its major metabolites L-dihydroxyphenylalanine, dopamine and 3,4-dihydroxyphenylacetic acid by HPLC with electrochemical detection 43
Diltiazem
Solid-phase extraction of diltiazem and its metabolites from plasma prior to HPLC 542
N-[2-(Dimethylamino)ethyl]acridine-4-carboxamide
Quantitation of the antitumour agent N-[2-(dimethylamino)ethyl]acridine-4-carboxamide in plasma by HPLC 385
2-(2,2-Diphenylcyclopropyl)-1*H*-imidazole
Quantification of cibenzoline and its imidazole metabolite by HPLC in human serum 267
Dopamine
Method for the determination of γ-L-glutamyl-L-dihydroxyphenylalanine and its major metabolites L-dihydroxyphenylalanine, dopamine and 3,4-dihydroxyphen-

ylacetic acid by HPLC with electrochemical detection 43

Dopamine
Reversed-phase LC method with amperometric detection for the determination of the dopamine agonist 2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin (N-0437) in human plasma and urine 464

Drug-free plasma profiles
Evaluation of drug-free plasma profiles by HPLC following on-line solid-phase extraction 277

Dyes
Analysis of xanthene dyes by reversed-phase HPLC on a polymeric column followed by characterization with a diode array detector 155

Felodipine
Enantioselective determination of felodipine and other chiral dihydropyridine calcium entry blockers in human plasma 343

*S*-Flunoxaprofen
Fluorescence assay for small peptides and amino acids: HPLC determination of selected substrates using activated *S*-flunoxaprofen as a chiral derivatizing agent 55

5-Fluorouracil
Stability-indicating HPLC method for the simultaneous determination of cisplatin and 5-fluorouracil in 0.9% sodium chloride for injection 516

Furazolidone
HPLC determination of furazolidone in Atlantic salmon (*Salmo salar*) tissue 495

Glucose utilization inhibitor
Isolation of an additional inhibitor of glucose utilization in renal insufficiency: pseudouridine 178

γ-L-Glutamyl-L-dihydroxypenylalanine
Method for the determination of γ-L-glutamyl-L-dihydroxyphenylalanine and its major metabolites L-dihydroxylphenylalanine, dopamine and 3,4-dihydroxyphenylacetic acid by HPLC with electrochemical detection 43

Glycopeptides
Rapid analysis of membrane glycopeptides by gel permeation HPLC 65

Guanine
Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91

Guanosine
Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91

Histones
Reactivity and adduct formation of a polyaromatic hydrocarbon, 7-bromomethylbenz[*a*]anthracene, with chromatin histone proteins 25

Human serum albumin
HPLC determination of the binding of ceftriaxone to human serum albumin solution and albumin from diluted human serum 415

10-Hydroxyamitriptyline
Enantiomer analysis of *E*- and *Z*-10-hydroxyamitriptyline in human urine 163

10-Hydroxycarbazepine
Simple and rapid micro-analytical HPLC technique for the assay of oxcarbazepine and its primary active metabolite 10-hydroxycarbazepine 473

Hydroxydiclofenac
HPLC method for the determination of diclofenac and its hydroxy metabolites in human plasma and urine 487

5-Hydroxyindoleacetic acid
Determination of six indolic compounds, including melatonin, in rat pineal using HPLC with serial fluorimetric–electrochemical detection 111

Hydroxymedifoxamine
Measurement of medifoxamine metabolites in urine by HPLC 531

5-Hydroxytryptophan
Determination of six indolic compounds, including melatonin, in rat pineal using HPLC with serial fluorimetric–electrochemical detection 111

Hypoxanthine
Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91

Ibuprofen
HPLC determination of the stereoisomeric metabolites of ibuprofen 395

Indoles
Determination of monoamines and indoles in amniotic fluid HPLC–electrochemical detection 101
Indoles
Determination of six indolic compounds, including melatonin, in rat pineal using HPLC with serial fluorimetric–electrochemical detection 111
Inosine
Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91
Inositol polyphosphates
Ion-pair chromatography of inositol polyphosphates with N-methylimipramine 143
Isethionic acid
Determination of isethionic acid by GC with flame photometric detection 172
Ivermectin
Improved HPLC method for quantitation of ivermectin in whole blood, serum or muscle tissue 250
γ-Linolenic acid
Quantitative assessment of γ-linolenic acid in human blood and plasma with capillary GC 447
L-toxin
Fast purification of *Clostridium sordellii* cytotoxin 357
Medifoxamine
Measurement of medifoxamine metabolites in urine by HPLC 531
Medifoxamine N-oxide
Measurement of medifoxamine metabolites in urine by HPLC 531
Melatonin
Determination of six indolic compounds, including melatonin, in rat pineal using HPLC with serial fluorimetric–electrochemical detection 111
Mepirodipine
Determination of mepirodipine in plasma by capillary column GC–negative-ion chemical ionization MS 191
N-Methylimipramine
Ion-pair chromatography of inositol polyphosphates with N-methylimipramine 143
Methylphenidate
Capillary GC–MS analysis of plasma methylphenidate 214
1-Methyl-1,2,3,4-tetrahydro-β-carboline
Highly sensitive method for the determination of 1-methyl-1,2,3,4-tetrahydro-β-carboline using combined capillary GC and negative-ion chemical ionization MS 1
Midazolam
Rapid, sensitive determination of human serum midazolam by HPLC 526
Molsidomine
Determination of the active metabolite of molsidomine in human plasma by reversed-phase HPLC 435
Monoamines
Determination of monoamines and indoles in amniotic fluid HPLC–electrochemical detection 101
Myelin basic protein
Rapid method based on reversed-phase HPLC for purification of human myelin basic protein and its thrombic and endoproteinase Lys-C peptides 79
N-0437
Reversed-phase LC method with amperometric detection for the determination of the dopamine agonist 2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin (N-0437) in human plasma and urine 464
Navelbine
HPLC determination of navelbine in human plasma and urine 258
Nicorandil
Rapid and simple determination of nicorandil in rat plasma using a solid-phase extraction column 509
Nucleobases
Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91
Nucleosides
Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91
Nucleotides
Measurement of adenine nucleotide levels with an adenine analyser as an index of freshness of porgy 305
Oxcarbazepine
Simple and rapid micro-analytical HPLC

technique for the assay of oxcarbazepine and its primary active metabolite 10-hydroxycarbazepine 473

Oxyphenbutazone
Estimation of the analgesic muscle relaxants chlorzoxazone and diazepam in human plasma by reversed-phase LC 407

Pelanserin
Determination of pelanserin in human plasma by HPLC with amperometric detection 501

Peptides
Fluorescence assay for small peptides and amino acids: HPLC determination of selected substrates using activated *S*-flunoxaprofen as a chiral derivatizing agent 55

Peptides
Rapid method based on reversed-phase HPLC for purification of human myelin basic protein and its thrombic and endoproteinase Lys-C peptides 79

Phencyclidine
Detection of low nanogram quantities of phencyclidine extracted from human urine. Preparation of an acetylated column packing material for use in GC with nitrogen–phosphorus detection 208

Plasma profiles, drug-free
Evaluation of drug-free plasma profiles by HPLC following on-line solid-phase extraction 277

Propofol
Simple and practical HPLC assay of propofol in human blood by phenyl column chromatography with electrochemical detection 537

2-(N-Propyl-N-2-thienylethylamino)-5-hydroxytetralin
Reversed-phase LC method with amperometric detection for the determination of the dopamine agonist 2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin (N-0437) in human plasma and urine 464

Proteins
Detection of the exercise-associated changes in serum proteins by two-dimensional electrophoresis under non-denaturing conditions 315

Pseudouridine
Isolation of an additional inhibitor of glucose utilization in renal insufficiency: pseudouridine 178

Purine nucleosides
Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91

Quinine
HPLC determination of quinine in plasma, whole blood and samples dried on filter paper 221

Scopolamine
Electon-impact ionization detection of scopolamine by GC–MS in rat plasma and brain 325

Serotonin
Determination of six indolic compounds, including melatonin, in rat pineal using HPLC with serial fluorimetric–electrochemical detection 111

Steroids
Analysis of corticosterone in rat urine by HPLC and fluorimetry using post-column reaction with sulphuric acid 453

Sulfamethoxazole
Ion-paired HPLC separation of trimethoprim, sulfamethoxazole and $N^4$-acetylsulfamethoxazole with solid-phase extraction 235

Thiodiglycol
Quantification of thiodiglycol in urine by electron ionization GC–MS 184

Trichloroacetic acid
Determination of chloral hydrate metabolism in adult and neonate biological fluids after single-dose administration 333

Trichloroethanol
Determination of chloral hydrate metabolism in adult and neonate biological fluids after single-dose administration 333

Trimethoprim
Ion-paired HPLC separation of trimethoprim, sulfamethoxazole and $N^4$-acetylsulfamethoxazole with solid-phase extraction 235

Trimethoprim
GC determination of trimethoprim in plasma during high-dose trimethoprim–sulfamethoxazole treatment for *Pneumocystis carinii* pneumonia 243

Tryptophan

Determination of six indolic compounds, including melatonin, in rat pineal using HPLC with serial fluorimetric–electrochemical detection 111

Verapamil

Measurement of verapamil in human plasma by reversed-phase HPLC using a short octyl column 480

Xanthene dyes

Analysis of xanthene dyes by reversed-phase HPLC on a polymeric column followed by characterization with a diode array detector 155

Xanthine

Determination of purine nucleosides and their bases by HPLC using co-immobilized enzyme reactors 91

Zidovudine

Analysis of 3′-azido-3′-deoxythymidine levels in tissues and milk by isocratic HPLC 371